



2017

Dielasma or Tunethyris? A Taxonomic Conundrum

Howard R. Feldman

Touro College, howard.feldman@touro.edu

Talia J. Belowich

Tova Braver

Sarah Laks

Follow this and additional works at: https://touroscholar.touro.edu/lcw_pubs

 Part of the [Paleontology Commons](#)

Recommended Citation

Feldman, H., Belowich, T., Braver, T., & Laks, S. (2017, March). Dielasma or Tunethyris? A taxonomic conundrum. Poster presented at the 2017 Geological Society of America Northeastern Section Meeting, Pittsburgh, PA.

This Poster is brought to you for free and open access by the Lander College for Women – The Anna Ruth and Mark Hasten School at Touro Scholar. It has been accepted for inclusion in Lander College for Women - The Anna Ruth and Mark Hasten School Publications and Research by an authorized administrator of Touro Scholar. For more information, please contact touro.scholar@touro.edu.



Tunethyris or Dielasma? a taxonomic conundrum





Feldman, Howard R.¹, Belowich, Talia J.¹, Braver, Tova² and Laks, Sarah²,
¹Division of Paleontology, American Museum of Natural History, 79th Street at Central Park West, New York,
NY 10024-5192, feldspar4@optonline.net; ²Biology Department, Touro College, New York, NY, 10023

Introduction

A new species of a dielasmid brachiopod from the Triassic Saharonim Formation of Makhtesh Ramon, southern Israel, was discovered in Anisian-Ladinian limestones that alternate with calcareous shales. The environment of deposition was an open shelf with normal salinity. The new species is homeomorphic with Paleozoic forms, but may not belong to the genus *Dielasma*; however, it may be a species of *Tunethyris*, a Triassic genus from Tunisia. If the Triassic specimens belong to the Paleozoic genus *Dielasma*, an important new Lazarus genus is added to the systematic literature. However, if the specimens belong to the Triassic *Tunethyris*, there are important paleobiogeographic considerations to be noted, such as the significance of the enigmatic Ladinian crisis and the influence of the Hispanic Corridor, an immature seaway connecting the western Tethys with eastern Panthalassa that may have been open sporadically during the Late Triassic and affected brachiopod distribution.

Tunethyris blodgetti n. sp. on left, *T. punica* on right. Serial sections of each are below.



Conclusions

Points favoring the selection of *Tunethyris* are:






1. The orientation of the crura in the new species is slightly different from Paleozoic dielasmids, especially *D. elongatum* from the Permian of Germany.
2. The anatomy of *Tunethyris punica* from the Norian of Tunisia is very similar to the new species and the anterior commissure of both is sulcinate, a feature not found in Paleozoic forms.
3. The loop of the Israeli brachiopods is acuminate. In addition to other anatomical characters that support the erection of a new species, *Tunethyris punica* has a septalium supported by a median septum whereas the septalium in the brachiopods from Makhtesh Ramon rests on the valve floor.

Acknowledgements

The fossils were collected during Feldman's tenure as a Visiting Scientist at the Geological Survey of Israel in Jerusalem. Thanks to Daniel Levin (USNM) for providing access to the collections. We are grateful to the following for discussions and an exchange of ideas related to the stratigraphy and depositional environments of the Triassic rocks in the Negev: Chaim Benjamini (Ben Gurion University, Beersheva, Israel), Hugo Bucher, Université Claude Bernard, Lyon, France), and Francis Hirsch (Naruto University of Education, Naruto, Japan). Support from the National Geographic Society is greatly appreciated.

References

- Calzada, S., B. Peybernes, and M.B. Youseff. 1994. *Tunethyris*, un nouveau genre de Brachiopode du Trias de Tunisie centrale. *Revue de Paléobiologie*, 13:117-124.
- Feldman, H. R., 2002. A new species of *Coenothyris* (Brachiopoda) from the Triassic (Ladinian) of Israel *Journal of Paleontology*, 76:34-42.
- Feldman, H. R., 2013. *Menathyris wilsoni* (Brachiopoda), a new genus and species from the Middle Triassic (Ladinian) of southern Israel. *Annales Societatis Geologorum Poloniae*, 83:81-85.
- Hirsch, F., 1972. Middle Triassic conodonts from Israel, southern France and Spain. *Mitteilungen Gesellschaft Geologie Bergbaustudenten*, 21:811-828.
- Stehli, F. G., 1956. *Dielasma* and its external homeomorph *Beecheria*. *Journal of Paleontology*, 30:299-302.
- Zak, I. 1963. Remarks on the stratigraphy and tectonics of the Triassic of Makhtesh Ramon. *Israel Journal of Earth Sciences*, 12:87-89



Paleozoic dielasmids

Top left, satellite view of eastern Mediterranean; circle denotes Makhtesh Ramon, Israel. Top right, stratigraphic column of southern Israel; bottom, arrow points to collecting locality.