

3-24-2016

Acute Promyelocytic Leukemia Co-Existing with JAK2 V617F Positive Myeloproliferative Neoplasm: A Case Report

Aleksandra Mamorska-Dyga
New York Medical College

Jingjing Wu

Pallavi Khattar
New York Medical College

Faisal Ronny
New York Medical College

Humayun Islam
New York Medical College

See next page for additional authors

Follow this and additional works at: https://touro scholar.touro.edu/nymc_fac_pubs



Part of the [Oncology Commons](#), and the [Pathology Commons](#)

Recommended Citation

Mamorska-Dyga, A., Wu, J., Khattar, P., Ronny, F. M., Islam, H., Seiter, K., & Liu, D. (2016). Acute promyelocytic leukemia co-existing with JAK2 V617F positive myeloproliferative neoplasm: A case report. *Stem Cell Investigation*, 3, 8. doi:10.21037/sci.2016.03.02

This Article is brought to you for free and open access by the Faculty at Touro Scholar. It has been accepted for inclusion in NYMC Faculty Publications by an authorized administrator of Touro Scholar. For more information, please contact touro.scholar@touro.edu.

Authors

Aleksandra Mamorska-Dyga, Jingjing Wu, Pallavi Khattar, Faisal Ronny, Humayun Islam, Karen Seiter, and Delong Liu

Acute promyelocytic leukemia co-existing with *JAK2* V617F positive myeloproliferative neoplasm: a case report

Aleksandra Mamorska-Dyga¹, Jingjing Wu^{1,2}, Pallavi Khattar³, Faisal M. H. Ronny³, Humayun Islam³, Karen Seiter¹, Delong Liu¹

¹Department of Medicine, Westchester Medical Center and New York Medical College, Valhalla, NY 10595, USA; ²Department of Oncology, The First Affiliated Hospital of Zhengzhou University, Zhengzhou 450000, China; ³Department of Pathology, Westchester Medical Center and New York Medical College, Valhalla, NY 10595, USA

Correspondence to: Delong Liu, MD, PhD. Professor of Medicine, Department of Medicine, Westchester Medical Center and New York Medical College, Valhalla, NY 10595, USA. Email: Delong_Liu@nymc.edu.

Abstract: The V617F mutation of Janus-associated kinase 2 (*JAK2*) is commonly seen in myeloproliferative neoplasms (MPN). Transformation of *JAK2* positive MPNs to acute leukemia has been reported. We here report a case of acute promyelocytic leukemia which was later confirmed to have a co-existing *JAK2* V617F positive MPN. In addition, the patient was found to have *FLT3*-TKD mutation, which, together with *PML/RARα*, could play a role in the MPN transformation to APL.

Keywords: Acute myelogenous leukemia (APL); myeloproliferative neoplasms (MPN); *PML/RARα*; Janus-associated kinase 2 (*JAK2*)

Received: 08 February 2016; Accepted: 09 March 2016; Published: 24 March 2016.

doi: 10.21037/sci.2016.03.02

View this article at: <http://dx.doi.org/10.21037/sci.2016.03.02>

Introduction

The Janus-associated kinase 2 (*JAK2*) V617F mutation is detected in over 95% of patients with polycythemia vera (PV) and in about 50% of cases of essential thrombocythemia (ET) and primary myelofibrosis (PMF) (1-3). Transformation of myeloproliferative neoplasms (MPN) into acute myelogenous leukemia (AML) is a well-studied and reported phenomenon (2,4,5). Both transformation of ET to AML (6,7), as well as a *JAK2* V617F mutation in *de novo* AML are very rare (8). Amongst all subtypes of AML originating from MPN, acute promyelocytic leukemia (APL) is extremely rare. In the English literature there have been only 9 such cases reported to date (1,7,9-14). In the same literature search we have not found any case of APL with concurrent diagnosis of MPN. Herein, we present the case of a young male with new onset APL and *JAK2* positive MPN.

Case presentation

A young male (<40 years old) was referred to our hospital

for gingival bleeding, pancytopenia and fever. The patient had previously presented with fever, sore throat, and tonsillar enlargement and had been treated empirically for pharyngitis with antibiotics. He also reported gingival bleeding and several episodes of epistaxis as well as dark stools in the few weeks preceding admission. Physical examination was significant for gingival bleeding and enlarged tonsils. Abdominal ultrasonography revealed splenomegaly with longest diameter of 14.3 cm. A complete blood count on admission was remarkable for a white blood count (WBC) of $2.1 \times 10^9/L$, hemoglobin (Hgb) of 9.8 g/dL, and platelet count of $11 \times 10^9/L$. The peripheral blood smear revealed 10% blasts and 42% promyelocytes. Bone marrow biopsy done on admission showed sheets of immature atypical myeloid cells, comprising more than 90% of the marrow cellularity. Those cells showed moderate to abundant cytoplasm, irregular nuclei (some of which were indented or bilobed), smudged chromatin and prominent nucleoli. Maturing myeloid elements including segmented neutrophils were not seen; megakaryocytes and erythroid precursors were markedly decreased. No reticulin fibrosis

Table 1 Cytogenetic and molecular profiles in the course of leukemia treatment

Date	t(15;17)	PML/RARa	JAK2 V617F (%)	FLT3 ITD	FLT3 TKD
7/19/2015	Positive [20/20]	Positive	34.00	Negative	Positive
8/17/2015	Negative	Weak positive	25.60	Negative	Negative
11/17/2015	ND	Negative	24.11	ND	ND

ITD, internal tandem duplication; TKD, tyrosine kinase domain mutation at D835; ND, not done.

Table 2 Molecular profiling by next generation sequencing of the bone marrow specimen

Profile	Genes	Result
MPN	<i>BCR/ABL, MPL, CALR, CSF3R, SETBP1</i>	–
MDS	<i>ASXL1, ETV6, EZH2, RUNX1, TP53</i>	–
AML		
RNA splicing	<i>SF3B1, SRSF2, U2AF1, ZRSR2</i>	–
Epigenetic	<i>ASXL1, DNMT3A, EZH2, IDH1/2, SETBP1, TET2</i>	–
Transcription factors	<i>CEBPa, ETV6, GATA2, PHF6, RUNX1, WT1</i>	–
Cohesion complex	<i>RAD1, SMC1A, SMC3, STAG2</i>	–
Activated signaling	<i>BRAF, CBL, KIT, KRAS, MPL, NF1, NRAS, PDGFa/b, PTPN11, STAT3, STAT5B</i>	–
Other	<i>BCOR, CALR, CSF3R, NPM1, TP53</i>	–

–, negative. MPN, myeloproliferative neoplasm; MDS, myelodysplastic syndrome; AML, acute myeloid leukemia.

was seen. Karyotyping revealed t(15;17)(q24;q21) in all 20 metaphases analyzed. Molecular studies were positive for PML/RARa, FLT3 TKD and *JAK2* V617F mutations (Table 1). Additional molecular studies for genes associated with MPN, MDS, and AML were negative (Table 2) (15). The patient was immediately started on all-trans retinoic acid (ATRA). Arsenic trioxide (ATO) was added on day 10 according to the reported regimen for clinically low-risk APL (16). The hospital course was complicated by differentiation syndrome with a high WBC (Figure 1A) and pleural and pericardial effusions for which a short course of dexamethasone was given. Surprisingly, a rapid increase in platelet count was observed during count recovery, with values reaching as high as 1,700×10⁹/L (Figure 1B). Bone marrow biopsy at this point showed increased reticulin fibrosis, a left shift in myeloid lineage cells with dysplastic and an increased number of megakaryocytes. The morphology overall was reported to be consistent with MPN (PMF/ET). A FISH panel for myelodysplasia was negative and a chromosome study revealed a normal karyotype: 46 XY. Molecular studies were negative for the *FLT3* TKD, but remained positive for the *JAK2* V617F mutation. The *PML/RARa* was still detectable by PCR post-induction therapy with ATRA and ATO. The patient was started on consolidation

chemotherapy with cytarabine and idarubicin for a total of two cycles. He continued ATRA throughout consolidation. The patient became PCR negative for *PML/RARa* after completion of two cycles of consolidation chemotherapy, however the *JAK2* mutation remained positive (Table 1). At that point the patient had a normal WBC but a high platelet count, with platelets as high as 949×10⁹/L. Since the patient presented with leukemia transformation, peak counts of platelets were higher than 1,500×10⁹/L, and bone marrow biopsy revealed dysplastic megakaryocytes as well as reticulin fibrosis, we believe that this patient might more likely have PMF than ET, and had high risk PMF/ET, even though he was younger than 40 (17). Therefore, low dose aspirin (81 mg) and hydroxyurea 1,000 mg daily were given for the PMF/ET. Interferon and ruxolitinib were discussed but clinically impractical at the time. The patient was also placed on maintenance treatment for APL with ATRA every 3 months.

Discussion

Both ET and PV may progress to myelofibrosis, and all MPNs may evolve into AML. The most common to transform in this way is PMF, occurring in about 15%




Figure 1 White blood counts and platelet counts during induction and consolidation chemotherapy. (A) The high WBC was observed during ATRA induction therapy. (B) Rapid increases in platelet count were seen during the recovery phase post chemotherapy.

Table 3 Case reports on transformation of myeloproliferative neoplasms to acute promyelocytic leukemia

Underlying MPN	Prior treatment	Years since diagnosis	References
ET→PMF	Hydroxyurea, aspirin, ruxolitinib	16.0→2.0	(1)
ET	None	9.3	(7)
ET	Uracil, mustard	3.9	(13)
ET	Busulfan	5.2	(14)
ET	Hydroxyurea	1.7	(11)
ET	Hydroxyurea, warfarin	4.0	(12)
PV	Phlebotomy	2.0	(12)
PV	Phlebotomy	9.3	(10)
PMF	Hydroxyurea	6.8	(9)

MPN, myeloproliferative neoplasm; ET, essential thrombocythemia; PMF, primary myelofibrosis; PV, polycythemia vera.

of cases (1,18). Such transformation happens even less frequently in patients with ET (6,7). In the majority of PV cases there are clones homozygous for *JAK2* V617F mutations, which are rarely found in ET (19,20). *JAK2* V617F mutations are very rare in *de novo* AML (8). Amongst all subtypes of AML originating from MPNs, APL has been reported the least frequently. Only 2 of the early case reports had molecularly documented *PML/RARα* mutation (7,12) (Table 3). The *JAK2* V617F mutation status was unknown in 8 of the cases (1), as they were reported prior to the 2005 discovery of the *JAK2* mutation (21,22). The association between APL and MPNs was thought to be due to promyelocytic blastic crisis of MPN, APL secondary

to cytoreductive therapy, and *de novo* APL (7). Braun *et al.* recently reported a case of APL with *PML/RARα* and *JAK2* V617F mutation (1). The authors postulated that the inflammatory response resulting from the chemokine release (CLL-2 and IL-8) from ATRA-treated APL cells was accentuated by inflammatory downstream signalling from the *JAK-2* mutation and that this led to severe differentiation syndrome. JAK-2 inhibition by ruxolitinib in the *in vitro* studies on NB-4 APL cells treated with ATRA did not affect CLL-2 and IL-8 levels, supporting the importance of *JAK2* in downstream activation as previously described (1,2). However, the role of JAK-2 inhibition in the management of differentiation syndrome has not been

clinically proven. In the patient presented by Braun *et al.* the diagnosis of MPN predated the diagnosis of APL. Our report presents for the first time a case of APL diagnosed concurrently with a MPN. It is highly possible that the APL clone remained dominant and masked the phenotype of MPN at the time of diagnosis. The MPN clone became dominant after the APL clone was suppressed. It remains unknown whether the APL clone arose from the MPN clone since we were not able to perform single cell genome analysis prior to the initiation of the APL therapy like in those cases reported in the literature (23–25). Since splenomegaly and *FLT3*-TKD were also present at the time of diagnosis, we hypothesize that the *JAK2* V617F mutated MPN clone (likely PMF) was present first and that additional mutations like *FLT3* and *PML/RARα* took place later and then led to the development of APL (26–32). It is of interest to point out that the case of APL transformed from a known diagnosis of ET with fibrosis also was found to have *FLT3*-TKD (D835 mutation) in addition to *JAK2* V617F mutation (1). These two cases of APL transformation from ET /PMF containing *FLT3*-TKD mutation make it likely that *FLT3*-TKD mutation plays a driver role for this transformation process involving *PML/RARα*.

Conclusions

This is the first case with molecular data showing co-existence of *PML/RARα*, *FLT3*-TKD, and a *JAK2* V617F mutation at the time of APL diagnosis. We hypothesize that the *JAK2* V617F mutated MPN clone was present first, and that additional mutations like *FLT3* and *PML/RARα* took place later and then led to the development of APL.

Acknowledgements

J Wu is a recipient of the Henan Provincial Grant for Overseas Research for Young Leaders of Medical Technology (No. 2014041). In addition, J Wu also received grant support from the Natural Science Foundation of China (NSFC No. 81201793). The grants supported her research training at the Division of Hematology and Oncology, New York Medical College, USA.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

Informed Consent: Unfortunately due to special circumstance in this case, consent could not be obtained for publication. The authors ensured that there was no identifiable information in the case (i.e., no race and age were reported). We believe this case has scientific value and is important for clinical literature.

References

1. Braun TP, Maxson JE, Agarwal A, et al. Acute promyelocytic leukemia with *JAK2* V617F and severe differentiation syndrome. *Leuk Res Rep* 2014;4:8–11.
2. Kralovics R, Passamonti F, Buser AS, et al. A gain-of-function mutation of *JAK2* in myeloproliferative disorders. *N Engl J Med* 2005;352:1779–90.
3. Furqan M, Mukhi N, Lee B, et al. Dysregulation of *JAK*-*STAT* pathway in hematological malignancies and *JAK* inhibitors for clinical application. *Biomark Res* 2013;1:5.
4. Björkholm M, Derolf AR, Hultcrantz M, et al. Treatment-related risk factors for transformation to acute myeloid leukemia and myelodysplastic syndromes in myeloproliferative neoplasms. *J Clin Oncol* 2011;29:2410–5.
5. Nagai Y, Kawahara M1, Sugino N, et al. A case of minor *BCR-ABL1* positive acute lymphoblastic leukemia following essential thrombocythemia and originating from a clone distinct from that harboring the *JAK2*-V617F mutation. *Exp Hematol Oncol* 2014;3:6.
6. Andersson PO, Ridell B, Wadenvik H, et al. Leukemic transformation of essential thrombocythemia without previous cytoreductive treatment. *Ann Hematol* 2000;79:40–2.
7. Sato N, Furukawa T, Masuko M, et al. Acute promyelocytic leukemia developing in untreated essential thrombocythemia. *Am J Hematol* 2002;71:114–6.
8. Theocharides A, Boissinot M, Girodon F, et al. Leukemic blasts in transformed *JAK2*-V617F-positive myeloproliferative disorders are frequently negative for the *JAK2*-V617F mutation. *Blood* 2007;110:375–9.
9. Batlle M, Fernández-Avilés F, Ribera JM, et al. Acute promyelocytic leukemia in a patient with idiopathic myelofibrosis. *Leukemia* 1999;13:492–4.
10. Kajiguchi T, Simokawa T, Saito M, et al. Transformation of polycythemia vera to acute promyelocytic leukemia. *Int J Hematol* 2000;72:520–1.
11. Löfvenberg E, Nordenson I, Wahlin A. Cytogenetic abnormalities and leukemic transformation in hydroxyurea-treated patients with Philadelphia chromosome negative chronic myeloproliferative disease.

- Cancer Genet Cytogenet 1990;49:57-67.
12. Mollee PN, Taylor KM, Williams B, et al. Long-term molecular remission in promyelocytic transformation of myeloproliferative disease. *Leukemia* 1999;13:648-50.
 13. Ratti M, Ghio R, Casciaro S, et al. Acute promyelocytic leukaemia developing in essential thrombocythaemia: blastic crisis or secondary acute leukaemia? Case report. *Haematologica* 1984;69:330-5.
 14. Sessarego M, Defferrari R, Dejana AM, et al. Cytogenetic analysis in essential thrombocythemia at diagnosis and at transformation. A 12-year study. *Cancer Genet Cytogenet* 1989;43:57-65.
 15. Li B, Gale RP, Xiao Z. Molecular genetics of chronic neutrophilic leukemia, chronic myelomonocytic leukemia and atypical chronic myeloid leukemia. *J Hematol Oncol* 2014;7:93.
 16. Lo-Coco F, Avvisati G, Vignetti M, et al. Retinoic acid and arsenic trioxide for acute promyelocytic leukemia. *N Engl J Med* 2013;369:111-21.
 17. Beer PA, Erber WN, Campbell PJ, et al. How I treat essential thrombocythemia. *Blood* 2011;117:1472-82.
 18. Cervantes F. How I treat myelofibrosis. *Blood* 2014;124:2635-42.
 19. Vannucchi AM. How I treat polycythemia vera. *Blood* 2014;124:3212-20.
 20. Tefferi A, Guglielmelli P, Larson DR, et al. Long-term survival and blast transformation in molecularly annotated essential thrombocythemia, polycythemia vera, and myelofibrosis. *Blood* 2014;124:2507-13; quiz 2615.
 21. James C, Ugo V, Casadevall N, et al. A JAK2 mutation in myeloproliferative disorders: pathogenesis and therapeutic and scientific prospects. *Trends Mol Med* 2005;11:546-54.
 22. James C, Ugo V, Le Couédic JP, et al. A unique clonal JAK2 mutation leading to constitutive signalling causes polycythaemia vera. *Nature* 2005;434:1144-8.
 23. Xie M, Lu C, Wang J, et al. Age-related mutations associated with clonal hematopoietic expansion and malignancies. *Nat Med* 2014;20:1472-8.
 24. Engle EK, Fisher DA, Miller CA, et al. Clonal evolution revealed by whole genome sequencing in a case of primary myelofibrosis transformed to secondary acute myeloid leukemia. *Leukemia* 2015;29:869-76.
 25. Hughes AE, Magrini V, Demeter R, et al. Clonal architecture of secondary acute myeloid leukemia defined by single-cell sequencing. *PLoS Genet* 2014;10:e1004462.
 26. Guglielmelli P, Lasho TL, Rotunno G, et al. The number of prognostically detrimental mutations and prognosis in primary myelofibrosis: an international study of 797 patients. *Leukemia* 2014;28:1804-10.
 27. Albano F, Anelli L, Zagaria A, et al. SETBP1 and miR_4319 dysregulation in primary myelofibrosis progression to acute myeloid leukemia. *J Hematol Oncol* 2012;5:48.
 28. Ferrer-Marín F, Bellosillo B, Martínez-Avilés L, et al. Leukemic transformation driven by an ASXL1 mutation after a JAK2V617F-positive primary myelofibrosis: clonal evolution and hierarchy revealed by next-generation sequencing. *J Hematol Oncol* 2013;6:68.
 29. Gelsi-Boyer V, Brecqueville M, Devillier R, et al. Mutations in ASXL1 are associated with poor prognosis across the spectrum of malignant myeloid diseases. *J Hematol Oncol* 2012;5:12.
 30. Randhawa J, Ostojic A, Vrhovac R, et al. Splenomegaly in myelofibrosis--new options for therapy and the therapeutic potential of Janus kinase 2 inhibitors. *J Hematol Oncol* 2012;5:43.
 31. Tognon R, Gasparotto EP, Neves RP, et al. Deregulation of apoptosis-related genes is associated with PRV1 overexpression and JAK2 V617F allele burden in Essential Thrombocythemia and Myelofibrosis. *J Hematol Oncol* 2012;5:2.
 32. Cole CB, Verdoni AM, Ketkar S, et al. PML-RARA requires DNA methyltransferase 3A to initiate acute promyelocytic leukemia. *J Clin Invest* 2016;126:85-98.

doi: 10.21037/sci.2016.03.02

Cite this article as: Mamorska-Dyga A, Wu J, Khattar P, Ronny FM, Islam H, Seiter K, Liu D. Acute promyelocytic leukemia co-existing with JAK2 V617F positive myeloproliferative neoplasm: a case report. *Stem Cell Investig* 2016;3:8.