

9-1-2017

## Drug-induced Causes of Secondary Hypertension

Wilbert S. Aronow  
*New York Medical College*

Follow this and additional works at: [https://touroscholar.touro.edu/nymc\\_fac\\_pubs](https://touroscholar.touro.edu/nymc_fac_pubs)



Part of the [Cardiology Commons](#), and the [Medical Pharmacology Commons](#)

---

### Recommended Citation

Aronow, W. S. (2017). Drug-induced Causes of Secondary Hypertension. *Annals of Translational Medicine*, 5 (17), 349. <https://doi.org/10.21037/atm.2017.06.16>

This Editorial is brought to you for free and open access by the Faculty at Touro Scholar. It has been accepted for inclusion in NYMC Faculty Publications by an authorized administrator of Touro Scholar. For more information, please contact [touro.scholar@touro.edu](mailto:touro.scholar@touro.edu).

# Drug-induced causes of secondary hypertension

Wilbert S. Aronow

Department of Medicine, Division of Cardiology, Westchester Medical Center and New York Medical College, Valhalla, NY, USA

Correspondence to: Wilbert S. Aronow, MD, FACC, FAHA. Professor of Medicine, Cardiology Division, Westchester Medical Center and New York Medical College. Macy Pavilion, Room 141 Valhalla, NY 10595, USA. Email: WSAronow@aol.com.

Submitted Apr 26, 2017. Accepted for publication May 11, 2017.

doi: 10.21037/atm.2017.06.16

View this article at: <http://dx.doi.org/10.21037/atm.2017.06.16>

Resistant hypertension is a blood pressure that remains above the treatment goal despite use of optimal doses of three antihypertensive drugs of different classes including a diuretic (1). Patients with resistant hypertension should be screened for causes of secondary hypertension (2). Common causes of secondary hypertension include renal parenchymal disease (1), renovascular disease (3), drug-induced causes (4), pregnancy (5), primary aldosteronism (6), and obstructive sleep apnea (7). Uncommon causes of secondary hypertension include acromegaly, hyperthyroidism, hypothyroidism, hyperparathyroidism, Cushing syndrome, apparent mineralocorticoid excess, pheochromocytoma/paraganglioma, carcinoid syndrome, congenital adrenal hyperplasia, coarctation of the aorta, and neurological causes (2). This article will discuss some of the drug-induced causes of secondary hypertension.

Excessive dietary sodium intake contributes to resistant hypertension by directly increasing blood pressure and by blunting the efficacy of most classes of antihypertensive drugs (8). This effect is especially pronounced in salt-sensitive persons such as older persons, blacks, and those with chronic kidney disease (1). Sodium-containing antacids should be avoided. Excessive alcohol use raises blood pressure and can cause resistance to antihypertensive drug therapy (4,9). Alcohol use should be restricted to  $\leq 2$  drinks daily for men and  $\leq 1$  drink daily for women (9). There is no difference in risk between beer, wine, and liquor (9). A meta-analysis of five studies in persons with hypertension showed that ingestion of 200 to 300 mg of caffeine caused an increase in blood pressure of 8.1/5.7 mm Hg, and the increase in blood pressure lasted for  $\geq 3$  hours (10). Persons with uncontrolled hypertension should avoid caffeinated beverages. Nicotine, cocaine, methamphetamine, and other recreational drugs which increase blood pressure should

be avoided (2,9). Cocaine causes a severe increase in blood pressure, especially if the person is taking a beta blocker (4). Smoking cessation decreases systolic blood pressure (9).

Because of their widespread use, nonsteroidal anti-inflammatory drugs, aspirin, and acetaminophen are the commonest drugs in worsening control of blood pressure (1). Blood pressure is increased by both cyclooxygenase 1-inhibitors and by cyclooxygenase 2-inhibitors. Nonsteroidal anti-inflammatory drugs increase blood pressure by influencing prostaglandin production causing adverse renal effects (9). Nonsteroidal anti-inflammatory drugs also increase systemic vascular resistance by increased endothelin-1 synthesis and by altered arachidonic metabolism (9). In 9,411 patients aged 65 years and older, use of nonsteroidal anti-inflammatory drugs increased the risk of starting antihypertensive drug treatment 1.55 times if low daily doses were used, 1.64 times if medium daily doses were used, and 1.82 times if high daily doses were used (11). Nonsteroidal anti-inflammatory drugs also blunt the blood-pressure lowering of antihypertensive drugs including diuretics, angiotensin-converting enzyme inhibitors, angiotensin receptor blockers, and beta-adrenergic blockers (12,13). Nonsteroidal anti-inflammatory drugs should be avoided if possible in persons with hypertension and other analgesics used depending on the indication.

Amphetamine, methylphenidate, dextromethylphenidate, and dextroamphetamine used to treat attention-deficit/hyperactivity disorder in children and adolescents may cause an elevated blood pressure (14). Behavioral therapies should be considered for the treatment of this disorder. Antidepressants such as monoamine oxidase inhibitors, serotonin-norepinephrine reuptake inhibitors, and tricyclic antidepressants may cause an elevated blood pressure (4).

Persons taking monoamine oxidase inhibitors must avoid foods and beverages containing high levels of tyramine such as liver, alcohol beverages, and aged cheeses or they may have a hypertensive crisis. Atypical antipsychotics such as clozapine and olanzapine may also cause an elevated blood pressure (15). Their use should be limited when possible.

Sympathomimetics such as the decongestants containing phenylephrine or pseudoephedrine may cause an elevated blood pressure (2). Alternative therapies such as nasal saline, intranasal corticosteroids, and antihistamines should be considered. Appetite suppressants may cause an elevated blood pressure (2). Herbal supplements such as ephedra, St John's wort, and yohimbine should be avoided since they have no clinical efficacy and may cause an elevated blood pressure (2).

Systemic corticosteroids such as dexamethasone, fludrocortisone, methylprednisolone, prednisone, and prednisolone may cause an elevated blood pressure (4). The increase in blood pressure is dose dependent. Alternative ways of administration of these drugs such as inhalation or topical use should be considered. Mineralocorticoids including licorice, carbenoxolone, 9- $\alpha$  fluorocortisol, ketoconazole, and carbenoxolone may cause an elevated blood pressure (4,16). The increase in blood pressure caused by these drugs is dose-dependent and is associated with hypokalemia, metabolic alkalosis, and a reduction in plasma renin activity and aldosterone levels (4,9).

Estrogens, androgens, and oral contraceptives may cause an increase in blood pressure (2,4,9). Oral contraceptives should not be used in women with uncontrolled hypertension (4). Low-dose ethinyl estradiol (20–30 mcg) agents should be used (4). Alternative methods of birth control should be used.

Immunosuppressants may cause an increase in blood pressure (2,4,9,17,18). Meta-analyses of randomized trials have demonstrated that cyclosporine treatment is associated with a higher incidence of hypertension than is treatment with tacrolimus (17,18).

Chronic use of recombinant human erythropoietin may cause hypertension (19). Mechanisms for the erythropoietin-induced hypertension include increase in hematocrit and erythrocyte mass, increased sensitivity to endogenous vasopressors, dysregulation of production or responsiveness to endogenous vasodilatory factors, a direct vasopressor effect, and stimulation of vascular cell growth (19). A sympathomimetic hypertensive crisis may be induced by abruptly stopping clonidine or other sympatholytic drugs or by using cocaine, amphetamines,

phencyclidine, or monoamine oxidase inhibitors (20).

## Acknowledgements

None.

## Footnote

*Conflicts of Interest:* The author has no conflicts of interest to declare.

## References

1. Calhoun DA, Jones D, Textor S, et al. Resistant hypertension: diagnosis, evaluation, and treatment: a scientific statement from the American Heart Association Professional Education Committee of the Council for High Blood Pressure Research. *Hypertension* 2008;51:1403-19.
2. Chiong JR, Aronow WS, Khan IA, et al. Secondary hypertension: current diagnosis and treatment. *Int J Cardiol* 2008;124:6-21.
3. Aronow WS. Atherosclerotic renal artery stenosis. *Ann Transl Med* 2017;5:264.
4. Grossman E, Messerli FH. Drug-induced hypertension: an unappreciated cause of secondary hypertension. *Am J Med* 2012;125:14-22.
5. Aronow WS. Hypertensive disorders in pregnancy. *Ann Transl Med* 2017;5:266.
6. Monticone S, Burrello J, Tizzani D, et al. Prevalence and clinical manifestations of primary aldosteronism encountered in primary care practice. *J Am Coll Cardiol* 2017;69:1811-20.
7. Pedrosa RP, Drager LE, Gonzaga CC, et al. Obstructive sleep apnea. The most common cause of hypertension associated with resistant hypertension. *Hypertension* 2011;58:811-7.
8. Weinberger MH, Sohen SJ, Miller JZ, et al. Dietary sodium restriction as adjunctive treatment of hypertension. *JAMA* 1988;259:2561-5.
9. Aronow WS, Fleg JL, Pepine CJ, et al. ACCF/AHA 2011 expert consensus document on hypertension in the elderly: a report of the American College of Cardiology Foundation Task Force on Clinical Expert Consensus Documents. Developed in collaboration with the American Academy of Neurology, American Geriatrics Society, American Society for Preventive Cardiology, American Society of Hypertension, American Society of Nephrology,

- Association of Black Cardiologists, and European Society of Hypertension. *J Am Coll Cardiol* 2011;57:2037-114.
10. Mesas AE, Leon-Munoz LM, Rodriguez-Artalejo F, et al. The effect of coffee on blood pressure and cardiovascular disease in hypertensive individuals: a systematic review and meta-analysis. *Am J Clin Nutr* 2011;94:1113-26.
  11. Gurwitz JH, Avorn J, Bohn RL, et al. Initiation of antihypertensive treatment during nonsteroidal anti-inflammatory drug therapy. *JAMA* 1994;272:781-6.
  12. Radack KL, Deck CC, Bloomfield SS. Ibuprofen interferes with the efficacy of antihypertensive drugs. A randomized, double-blind, placebo-controlled trial of ibuprofen compared with acetaminophen. *Ann Intern Med* 1987;107:628-35.
  13. Conlin PR, Moore TJ, Swartz SL, et al. Effect of indomethacin on blood pressure lowering by captopril and losartan in hypertensive patients. *Hypertension* 2000;36:461-5.
  14. Cortese S, Holtmann M, Banaschewski T, et al. Practitioner review: current best practice in the management of adverse events during treatment with ADHD medications in children and adolescents. *J Child Psychol Psychiatry* 2013;54:227-46.
  15. Newcomer JW. Antipsychotic medications: metabolic and cardiovascular risk. *J Clin Psychiatry* 2007;68:8-13.
  16. Chobanian AV, Volicer L, Tiffet CP, et al. Mineralocorticoid-induced hypertension in patients with orthostatic hypotension. *N Engl J Med* 1979;301:68-73.
  17. Liu Y, Yang MS, Yuan JY. Immunosuppressant utilization and cardiovascular complications among Chinese patients after kidney transplantation: a systematic review and analysis. *Int Urol Nephrol* 2013;45:885-92.
  18. Xue W, Zhang Q, Xu Y, et al. Effect of tacrolimus and cyclosporine treatment on metabolic syndrome and cardiovascular risk factors after renal transplantation: a meta-analysis. *Chin Med J* 2014;127:2376-81.
  19. Vaziri ND. Mechanism of erythropoietin-induced hypertension. *Am J Kidney Dis* 1999;33:821-8.
  20. Aronow WS. Treatment of hypertensive emergencies. *Ann Transl Med* 2017;5:S5.

**Cite this article as:** Aronow WS. Drug-induced causes of secondary hypertension. *Ann Transl Med* 2017;5(17):349. doi: 10.21037/atm.2017.06.16