

9-1-2017

Reply to Aitken et al., "Should Piperacillin-Tazobactam Be Used as Definitive Therapy Against Enterobacteriaceae Harboring Inducible AmpC β -Lactamases?"

L Cheng

B Nelson

M Mehta

Qiuhi Shi

New York Medical College

A Gomez-Simmonds

See next page for additional authors

Follow this and additional works at: https://touro scholar.touro.edu/nymc_fac_pubs



Part of the [Medicine and Health Sciences Commons](#)


Recommended Citation

Cheng, L., Nelson, B., Mehta, M., Shi, Q., Gomez-Simmonds, A., & Uhlemann, A. (2017). Reply to Aitken et al., "Should Piperacillin-Tazobactam Be Used as Definitive Therapy Against Enterobacteriaceae Harboring Inducible AmpC β -Lactamases?". *Antimicrobial Agents and Chemotherapy*, 61 (9), e01163-17. <https://doi.org/10.1128/AAC.01163-17>

This Response or Comment is brought to you for free and open access by the Faculty at Touro Scholar. It has been accepted for inclusion in NYMC Faculty Publications by an authorized administrator of Touro Scholar. For more information, please contact touro.scholar@touro.edu.

Authors

L Cheng, B Nelson, M Mehta, Qiuhi Shi, A Gomez-Simmonds, and A Uhlemann



Reply to Aitken et al., “Should Piperacillin-Tazobactam Be Used as Definitive Therapy against *Enterobacteriaceae* Harboring Inducible AmpC β -Lactamases?”

Lucy Cheng,^a Brian C. Nelson,^b Monica Mehta,^b Qiuhiu Shi,^c
Angela Gomez-Simmonds,^a Anne-Catrin Uhlemann^a

Department of Medicine, Division of Infectious Diseases, Columbia University Medical Center, New York, New York, USA^a; New York-Presbyterian Hospital, New York, New York, USA^b; New York Medical College, Valhalla, New York, USA^c

We appreciate Aitken et al.’s careful review (1) of our recent paper assessing the use of piperacillin-tazobactam for the treatment of bloodstream infections caused by *Enterobacteriaceae* harboring AmpC β -lactamases (2). We respectfully disagree with their assertion that the numerical increase in mortality seen with the use of piperacillin-tazobactam compared to the mortalities of patients on carbapenems and/or cefepime in our and other studies suggests that inferior outcomes are to be expected with piperacillin-tazobactam.

Aitken et al. are correct to point out that the odds ratio (OR) for the 30-day survival, not mortality, of patients receiving piperacillin-tazobactam versus that of patients receiving cefepime or meropenem was 0.5 in the propensity-matched data set. Therefore, the OR for 30-day mortality was 2.0 (95% confidence interval, 0.5 to 7.7), representing 3 additional deaths in the piperacillin-tazobactam group. This result was not statistically significant ($P = 0.33$), indicating that we were unable to detect an association between the treatment regimen and the occurrence of mortality in this study. It is worth noting that the large confidence intervals reported here suggest a low level of the precision of the OR. Thus, it is inappropriate to draw a firm conclusion about the relative odds of mortality based on its numerical value alone. As we pointed out in our publication, propensity score matching lowered our ability to detect differences in outcomes. In the overall cohort, the OR for 30-day survival was 1.16, as reported, and the OR for 30-day mortality was 0.86 (95% confidence interval, 0.3 to 2.3; $P = 0.8$). It is worth noting that in creating the propensity score-matched cohort, fewer deaths were excluded from the piperacillin-tazobactam group (3/9) than from the carbapenem/cefepime group (6/9).

Aitken et al. also suggested that using multivariable logistic regression is preferred to propensity score matching for some data sets. We opted for propensity score matching to minimize differences between baseline characteristics of treatment groups. In addition, a low number of outcome events in our data set made it difficult to perform multivariate models. Multivariable analyses of the overall cohort were attempted by incorporating variables that were significantly different between treatment groups, but these models demonstrated poor fit. However, they also showed no association between treatment groups and mortality, supporting our findings from the propensity score-matched analysis.

The commenters further mention a paper by Harris and colleagues from 2017 with similar results (i.e., a numerically higher rate of recurrence of *Enterobacter* spp. bacteremia without a statistically significant difference) and a small patient population (3).

Citation Cheng L, Nelson BC, Mehta M, Shi Q, Gomez-Simmonds A, Uhlemann A-C. 2017. Reply to Aitken et al., “Should piperacillin-tazobactam be used as definitive therapy against *Enterobacteriaceae* harboring inducible AmpC β -lactamases?” *Antimicrob Agents Chemother* 61:e01163-17. <https://doi.org/10.1128/AAC.01163-17>.

Copyright © 2017 American Society for Microbiology. All Rights Reserved.

Address correspondence to Anne-Catrin Uhlemann, au2110@columbia.edu.

This is a response to a letter by Aitken et al. (<https://doi.org/10.1128/AAC.01160-17>).

We direct them to a much larger meta-analysis also conducted by Harris et al. in 2016 that included 11 studies and revealed no strong evidence to suggest the superiority of carbapenems to other treatment regimens, including β -lactamase/ β -lactamase inhibitors, in patients with bloodstream infections due to *Enterobacteriaceae* with AmpC β -lactamases (4). We welcome future studies of the efficacy of non-carbapenem-containing regimens, such as piperacillin-tazobactam, for the treatment of AmpC-harboring *Enterobacteriaceae*. As we discussed in our original paper, prospective studies and meta-analyses are needed to further delineate patient populations that might benefit from this treatment approach and to identify specific risk factors warranting the use of carbapenem or cefepime.

REFERENCES

1. Aitken SL, Foolad F, Mcdaneld PM. 2017. Should piperacillin-tazobactam be used as definitive therapy for *Enterobacteriaceae* harboring inducible AmpC β -lactamases? *Antimicrob Agents Chemother* 61:e01160-17. <https://doi.org/10.1128/AAC.01160-17>.
2. Cheng L, Nelson BC, Mehta M, Seval N, Park S, Giddins MJ, Shi Q, Whittier S, Gomez-Simmonds A, Uhlemann AC. Piperacillin-tazobactam versus other antibacterial agents for treatment of bloodstream infections due to AmpC β -lactamase-producing *Enterobacteriaceae*. *Antimicrob Agents Chemother* 61:e00276-17. <https://doi.org/10.1128/AAC.01163-17>.
3. Harris PNA, Peri AM, Pelecanos AM, Hughes CM, Paterson DL, Ferguson JK. 2017. Risk factors for relapse or persistence of bacteraemia caused by *Enterobacter* spp.: a case-control study. *Antimicrob Resist Infect Control* 6:14. <https://doi.org/10.1186/s13756-017-0177-0>.
4. Harris PN, Wei JY, Shen AW, Abdile AA, Paynter S, Huxley RR, Pandeya N, Doi Y, Huh K, O'Neal CS, Talbot TR, Paterson DL. 2016. Carbapenems versus alternative antibiotics for the treatment of bloodstream infections caused by *Enterobacter*, *Citrobacter* or *Serratia* species: a systematic review with meta-analysis. *J Antimicrob Chemother* 71:296–306. <https://doi.org/10.1093/jac/dkv346>.