

12-1-2018

Regional Tissue Oxygen Extraction and Severity of Anemia in Very Low Birth Weight Neonates: A Pilot NIRS Analysis

Jonathan Mintzer

Boriana Parvez
New York Medical College

Edmund F. La Gamma
New York Medical College

Follow this and additional works at: https://touro scholar.touro.edu/nymc_fac_pubs



Part of the [Medicine and Health Sciences Commons](#)

Recommended Citation

Mintzer, J., Parvez, B., & La Gamma, E. (2018). Regional Tissue Oxygen Extraction and Severity of Anemia in Very Low Birth Weight Neonates: A Pilot NIRS Analysis. *American Journal of Perinatology*, 35 (14), 1411-1418. <https://doi.org/10.1055/s-0038-1660458>

This Article is brought to you for free and open access by the Faculty at Touro Scholar. It has been accepted for inclusion in NYMC Faculty Publications by an authorized administrator of Touro Scholar. For more information, please contact touro.scholar@touro.edu.

Regional Tissue Oxygen Extraction and Severity of Anemia in Very Low Birth Weight Neonates: A Pilot NIRS Analysis

Jonathan P. Mintzer, MD,¹ Borianna Parvez, MD,² and Edmund F. La Gamma, MD²

¹Department of Pediatrics, Division of Neonatal-Perinatal Medicine

Stony Brook Children's Hospital, Stony Brook, NY, USA

²Department of Pediatrics, Division of Newborn Medicine

Maria Fareri Children's Hospital at Westchester Medical Center, Valhalla, NY, USA

Running Title: Hematocrit and Tissue Oxygen Extraction

Corresponding Author:

Jonathan P. Mintzer, MD (author of manuscript first draft)

Assistant Professor of Pediatrics, Stony Brook Children's Hospital

Department of Pediatrics, Division of Neonatal-Perinatal Medicine

HSC T11, Rm. 060, Stony Brook, New York 11794-8111

Telephone: 631 444 7653

Fax: 631 444 8968

Email: jonathan.mintzer@stonybrookmedicine.edu

Financial Support: This study was supported in part by an unrestricted equipment use grant from Somanetics (now Medtronic, Boulder, CO).

Disclosure: All authors have no financial relationships or conflicts of interest to disclose related to this study.

ABSTRACT

Objective: Anemia causes blood flow redistribution and altered tissue metabolic behavior to sustain homeostatic oxygen consumption. We hypothesized that anemia severity would correlate with increased regional fractional tissue oxygen extraction among premature neonates.

Study Design: Regional oxygen extraction was calculated using pulse oximetry and near-infrared spectroscopy data among neonates <1250g during their first ten postnatal days. Oxygen extraction was assessed for correlations with raw hematocrit levels and following grouping into hematocrit quartiles.

Results: Twenty-seven neonates with gestational age 27 ± 2 wk and birth weight 966 ± 181 g underwent 116 hematocrit determinations. Cerebral and flank oxygen extraction inversely correlated with hematocrit (cerebral $r = -0.527$, $p = 0.005$; flank $r = -0.485$, $p = 0.01$). Increased cerebral oxygen extraction was observed for the lowest three hematocrit quartiles (Q1 0.26 ± 0.08 , $p=0.004$; Q2 0.24 ± 0.09 , $p=0.01$; Q3 0.25 ± 0.09 , $p=0.03$; all compared to Q4 0.18 ± 0.10). Increased flank oxygen extraction occurred for the lowest two quartiles (Q1 0.36 ± 0.12 , $p<0.001$; Q2 0.35 ± 0.11 , $p<0.001$; compared to Q4 0.22 ± 0.13). Splanchnic oxygen extraction demonstrated no similar correlations.

Conclusion: Increases in tissue oxygen extraction may indicate early pathophysiologic responses to nascent anemia in premature neonates.

Keywords: Anemia, hematocrit, neonate, NIRS, oxygen extraction, regional oxygenation

INTRODUCTION

Anemia occurs commonly in very low birth weight (VLBW) neonates.¹ In the neonatal intensive care unit (NICU), transfusion requirements are most-often defined by serial hematocrit or hemoglobin determinations.² Currently, hematocrit and clinical thresholds for packed red blood cell (pRBC) transfusion remain controversial.²⁻⁴ What is agreed upon is that progressive anemia can create an oxygen delivery/consumption imbalance resulting in blood flow maldistribution with variations in local tissue oxygen extraction. In extreme situations, insufficient oxygen delivery results in tissue acidosis, previously illustrated in neonates by direct measurement using micro-pH electrodes, with associated systemic metabolic acidosis resulting from a shift to anaerobic metabolism.⁵ To maintain homeostatic stability among VLBW neonates, these observations portend the need for more continuous, physiologically-based assessments of oxygenation adequacy to meet consumption demands during progressive anemia.⁶

Numerous studies utilize near-infrared spectroscopy (NIRS) for noninvasive bedside monitoring of *regional tissue oxygenation* (rSO₂). Whereas pulse oximetry-derived oxygen saturation (SpO₂) estimates anticipated tissue oxygen *delivery* from arterialized vessels, NIRS provides oxygen saturation data from predominantly post-capillary hemoglobin sources.^{7,8} By simultaneously monitoring SpO₂ and tissue-specific rSO₂ via NIRS (e.g., cerebral, flank/renal, splanchnic), one can calculate a trendable estimate of fractional tissue oxygen extraction ($FTOE = (SpO_2 - \text{region-specific } rSO_2) / SpO_2$).⁹⁻¹¹ Identifying increases in tissue oxygen extraction could allow earlier identification of blood flow maldistribution or insufficient tissue oxygen delivery to meet metabolic demands before pathologic clinical sequelae develop.

Although previous reports have described effects of physiologic perturbations in oxygen delivery on tissue rSO₂ and FTOE,^{9,12-14} data are lacking in VLBW neonates regarding organ-specific oxygen extraction relationships relative to the magnitude of anemia.^{12,15} In this non-*a priori* pilot analysis, we sought to determine whether hematocrit values correlate with region-specific FTOE. Further, we aimed to investigate the tissue-specific oxygen extraction responses to varied degrees of anemia. We hypothesized that decreased hematocrit levels would be associated with progressively increased cerebral, flank, and splanchnic FTOE and that these changes in oxygenation behavior would demonstrate regional specificity.

METHODS

Study Design

This prospective observational cohort study was approved by the Institutional Review Board of New York Medical College and the Maria Fareri Children's Hospital as minimal risk. Data were acquired during a NIRS survey of VLBW neonates to evaluate region-specific baseline values and changes occurring with routine clinical practices over the first 10 postnatal days. Previous work related to this patient cohort and data set have evaluated booster pRBC transfusions,¹⁰ quiescent baseline variability,¹⁶ sodium bicarbonate corrections,¹⁷ and effects of umbilical arterial blood drawing.¹⁸ This hypothesis-generating pilot analysis was conducted *post hoc* to evaluate the relationship between hematocrit level and region-specific FTOE. The current objective was to determine whether FTOE is sensitive enough for use in a larger randomized study designed to examine the graded contributions of various factors affecting oxygen delivery/consumption balance in VLBW neonates.

Patient Population

Subjects were admitted to the 50-bed, Level IV Regional Perinatal Center at Maria Fareri Children's Hospital of Westchester Medical Center from a catchment area of 23,000 annual births. Following parental consent, VLBW neonates < 1250 grams birth weight were enrolled within the first 72 postnatal hours and were monitored for an additional seven days each. Newborns with birth weights less than 500 grams, major congenital/chromosomal anomalies, congenital heart disease, five minute Apgar score <3, not expected to survive for the study duration, or with extremely immature skin were excluded.

Monitoring

After delivery and initial management, continuous cerebral, flank, and splanchnic NIRS monitoring was initiated using the INVOS 5100C Cerebral/Somatic Oximeter with OxyAlert NIR Sensors – Infant Model IS [Neonatal] (Somanetics, Troy, MI, now Medtronic, Boulder, CO). This device estimates the regional oxygenation status of underlying venous, arterial, and capillary hemoglobin sources using an assumed 75:20:5 respective ratio,¹⁹ at a manufacturer-reported signal depth of 1.5cm beneath the monitoring sensor.²⁰ Regional tissue oxygen saturation (rSO₂) is displayed as a proportion of oxygenated to total hemoglobin [oxyhemoglobin/(oxyhemoglobin + deoxyhemoglobin)].⁸ Consistent with numerous previous studies, NIRS sensor placement was as follows:²¹

Cerebral: Transverse across forehead

Flank/Renal: Longitudinally in right paraspinal region

Splanchnic: Transverse in infraumbilical position or obliquely from left lower

abdominal quadrant toward left flank if space limited

To protect fragile neonatal skin, NIRS monitoring sensors were placed atop Mepitel gel-impregnated gauze strips (Mölnlycke Health Care, Göteborg, Sweden). This technique was confirmed by Somanetics (now Medtronic) and our team as not affecting NIRS signal integrity while conferring adequate skin protection.²¹ Monitoring sensors were removed daily for a 30 minute “*skin resting period*” to provide ongoing assessment of skin integrity.

NICU Care / Hematocrit Measurements

All NICU care and decisions regarding hematocrit determinations were rendered by attending neonatologists. Hematocrits were obtained as a component of complete blood counts using a Coulter Counter (Beckman Coulter, Brea, CA). Typically, subjects had a hematocrit drawn upon NICU admission; further hematocrit determinations were based on attending discretion. No prospective decisions, including additional laboratory testing, were made based on NIRS results. Given previously published work from this data set regarding the effects of booster pRBC transfusions on FTOE,¹⁰ post-transfusion hematocrits were excluded from analysis in order to evaluate the relationship between hematocrit and FTOE to potentially determine transfusion thresholds for a future randomized trial.

Data Collection and Analysis

Antenatal, birth history, and hematocrit data were abstracted from subjects’ medical records. Raw NIRS data were collected every 6 seconds; the most frequent data sampling setting offered by the INVOS 5100C device. For each hematocrit determination, corresponding time-synchronized pulse oximetry and cerebral, flank, and

splanchnic NIRS data were collected over 15 minute epochs to allow for simultaneous FTOE calculations $((\text{SpO}_2 - \text{region-specific rSO}_2) / \text{SpO}_2)$. Based on previous variability analysis, this epoch length was selected to minimize the signal-to-noise ratio resulting from differences in region-specific baseline variability.¹⁶

Statistics

Comparisons between cerebral, flank, and splanchnic rSO₂ and FTOE were performed using ANOVA following confirmation of normal distributions. Scatterplots were created comparing hematocrits with organ-specific FTOE values. Pearson (r) calculations were then performed to evaluate for correlations between hematocrits and region-specific rSO₂ and FTOE values. Given multiple non-independent measures, Pearson (r) analyses were additionally conducted on both pooled raw data and averaged individual-patient data.²² Analyses of NIRS-derived data were also performed following organization of hematocrit data into quartiles. Comparisons across quartiles were performed via ANOVA (or ANOVA on ranks) with Dunnett (or Dunn's, if ranks used) post-hoc test using the highest hematocrit quartile (Q4) as baseline (normal/ideal hematocrit). Finally, multivariate logistic regression modeling was performed to evaluate the regional FTOE effects of hypotension requiring inotropic support; hemodynamically-significant patent ductus arteriosus; and intraventricular hemorrhage. Statistical significance was designated as a P value of < 0.05. All statistical analyses were performed using SigmaPlot v12.0 (Aspire Software International, Ashburn, VA).

RESULTS

Twenty-seven VLBW neonates underwent continuous NIRS monitoring between 6 and 221 hours of life. A total of 116 hematocrit determinations were performed and

verified as not preceded by pRBC transfusions during the prior 72 hours. Gestational age was 27 ± 2 wk (mean \pm SD) and birth weight was 966 ± 181 g. Cesarean deliveries occurred in 59% (16/27) of subjects, delivery room intubation in 56% (15/27), and 15% (4/27) had 5-minute Apgar scores less than 7. During the seven days of NIRS monitoring, 11% (3/27) of subjects received short courses of dopamine for hypotension, 41% (11/27) were diagnosed with patent ductus arteriosus, and 26% (7/27) demonstrated intraventricular hemorrhage on routine head ultrasound. Hematocrit and corresponding vital sign data are presented in Table 1.

Cerebral, flank, and splanchnic individual-hematocrit raw data scatterplots with overall data regression lines and Pearson r correlations with 95% confidence intervals are displayed in Figure 1. As individual subjects contributed between 1 and 10 hematocrit values to this data set, an additional analysis was conducted using averaged individual-subject data to account for non-independent/repeated measures,²² with no changes in correlations noted following this sub-analysis. Additional correlational analyses demonstrated direct relationships between hematocrit and raw cerebral and flank rSO_2 data (cerebral $r = 0.418$ and flank $r = 0.436$, both $p < 0.01$). No correlation was observed between hematocrit and raw splanchnic rSO_2 data ($r = 0.137$, $p = 0.15$).

Hematocrit quartiles (Figure 2) were subsequently analyzed for correlations with NIRS-derived data (Table 2 and Figure 3-4). Across all measurements, cerebral rSO_2 demonstrated the highest and least variable signal, with flank providing intermediate findings, and splanchnic oxygenation uniformly showing the lowest and most variable rSO_2 data ($p < 0.05$ for comparisons of both value and variability) as previously shown in other studies utilizing portions of this data set.^{9,15,16} Calculated FTOE was smallest and

least variable for the cerebral vascular bed, intermediate for the flank site, and largest with the greatest variability for splanchnic measurements ($p < 0.05$ for comparisons of both value and variability).

Relationships between hematocrit quartiles and organ-specific rSO_2 and FTOE are shown in Table 2 and Figures 3-4. Cerebral rSO_2 was diminished, with corresponding elevated FTOE, for the three lowest hematocrit quartiles compared to the highest quartile. A more graded relationship was observed between hematocrit quartile and renal rSO_2 and FTOE. No statistically significant correlations were noted across hematocrit quartiles for splanchnic NIRS-derived data.

On multivariate logistic regression, the effect of hematocrit on cerebral and flank FTOE trends remained unaltered in the presence of single or multiple factors potentially affecting organ perfusion. Interestingly, hematocrit demonstrated a significant correlation with splanchnic FTOE only in the presence of other risk factors, and particularly patent ductus arteriosus. Multivariate logistic regression modeling controlling for hypotension requiring inotropic support; presence of hemodynamically-significant patent ductus arteriosus; and presence of intraventricular hemorrhage are displayed in Table 3.

DISCUSSION

This report demonstrates that in VLBW neonates during the first ten postnatal days, cerebral and flank FTOE are inversely correlated with hematocrit levels. No similar correlation was observed for the splanchnic monitoring site. Across all hematocrit values, oxygen extraction was lowest in the brain, followed by flank and splanchnic vascular beds, consistent with neuroprotective maintenance of cerebral oxygen delivery.⁹ These findings suggest potential practical utility for NIRS-based trend monitoring as an

adjunctive, proportional measure of regional oxygen delivery/consumption balance among premature neonates. As homeostatic perturbations are most observable during later stages of oxygen privation,²³ these data may contribute to earlier appreciation of minimally symptomatic anemia prior to clinically apparent distress.

Anemia, both physiologic and iatrogenic, remains a common diagnosis in the NICU setting.^{1,4} Unfortunately, in VLBW neonates, the need for frequent phlebotomy during the first several postnatal days contributes to anemia itself.¹⁰ While strategies to reduce phlebotomy have been employed, relatively frequent phlebotomy is common among the smallest, most premature neonates with the lowest total blood volume and red cell mass.^{2,4} Thus, techniques to noninvasively assess oxygen delivery/consumption balance in VLBW neonates before clinical symptoms occur represent a potential advancement.

As anemia reduces blood oxygen-carrying capacity, our data confirm the hypothesis that individual tissues demonstrate enhanced oxygen extraction at lower hematocrit levels. Other studies have also employed NIRS to investigate anemia in premature neonates. Common to these reports are increases in region-specific rSO₂ (or decreases in FTOE) following transfusion, whether provided for symptomatic anemia^{9,12-14,24,25} or as a booster transfusion for phlebotomy-related blood losses.¹⁰ However, none of these studies included correlational analyses to ascertain regional rSO₂ or FTOE values based solely on anemia severity.

On logistic regression modeling, decreased hematocrit remained correlated with elevated cerebral and flank FTOE even when other factors were associated with enhanced tissue oxygen extraction. Interestingly, a relationship between hematocrit and splanchnic

oxygen extraction was observed only when other perfusion-affecting comorbidities existed, particularly patent ductus arteriosus. Given the inherent variability of splanchnic monitoring,¹⁶ it currently appears that this correlation only becomes “visible” during enhanced stages of oxygen privation. A larger future study encompassing a wider range of hematocrit values is required to validate this interesting result.

In this data set, while NIRS sensor placement is consistent with previous neonatal studies, concerns remain regarding actual tissue sampling, as regional monitoring can be affected by neighboring tissues’ perfusion. For example, though flank monitoring is assumed to estimate renal oxygenation, one cannot be certain of actual renal monitoring without imaging for kidney position. Additionally, as previously reported, excessive splanchnic variability results in significant limitations in interpreting splanchnic FTOE data.^{9,21} Moreover, some reports have indicated enhanced splanchnic variability as a risk factor for various outcomes, including transfusion-related acute gut injury.²⁶ While data averaging techniques may correct for baseline splanchnic variability, substantial uncertainty remains regarding the utility of splanchnic NIRS monitoring for clinical decision-making. Further studies, including innovative data smoothing methods, are needed to properly evaluate the possible clinical utility of splanchnic monitoring.

Whereas rSO₂ monitoring is currently used for trend analysis in specific care settings (i.e. during surgery), NIRS monitoring is not yet routinely employed in the NICU. Should continuous rSO₂ and FTOE monitoring become available at the bedside and a maximal FTOE “*zone of comfort*” established, this adjunctive, noninvasive measure could reduce dependence on empiric hematocrit thresholds for transfusions. Importantly, establishing rSO₂ and/or FTOE trends is likely more clinically relevant than absolute

values as individual patients may produce their own baselines to monitor for concerning trends (e.g., decreasing rSO₂ or increasing FTOE). Given the numerous factors which affect oxygen delivery/consumption balance in neonates, this noninvasive monitoring strategy may allow enhanced discretion for consideration of invasive bedside testing practices, including frequent blood sampling. Though larger, rigorous prospective studies are required, the NIRS modality shows promise toward providing a more individualized, physiologically-based approach for managing oxygen delivery/consumption balance among vulnerable neonates.

This study's primary strength is its proof of concept utilizing continuous NIRS monitoring with a reasonably large number and wide range of hematocrit determinations obtained from VLBW neonates during the first 10 postnatal days. However, several limitations affect the utility and generalizability of the results. Primarily, this hypothesis-generating analysis utilized a small patient sample size and non-*a priori* study design. Indeed, this report utilizes data that were collected during a prospective, normative NIRS survey among VLBW neonates not specifically designed to evaluate the relationship between hematocrit and regional FTOE. Moreover, given the large number of factors potentially affecting FTOE (e.g. inotropic support,²⁷ patent ductus arteriosus,²⁸ intraventricular hemorrhage,²⁹ enteral feeding status,³⁰ and others), important questions remain regarding the relative contribution of these parameters to tissue oxygen extraction. In addition, subtle yet significant differences in systolic and diastolic blood pressure and oxygen saturation between hematocrit quartile groups may have affected observed oxygen extraction behavior. Finally, hematocrits obtained based on clinical discretion may have inadvertently introduced a confounding selection bias based on clinical practice

variability. A larger prospective study, designed to account for the numerous confounding variables affecting oxygen delivery/consumption balance, is required to further develop FTOE trending in the management of anemia in VLBW neonates.^{12,15}

In summary, in this pilot study of the relationship between hematocrit and cerebral, flank, and splanchnic FTOE, hematocrit level was inversely correlated with cerebral and flank oxygen extraction in VLBW neonates. A similar relationship was not observed for the splanchnic vascular bed, except in the presence of hemodynamically-significant patent ductus arteriosus. Further study is required to examine the graded effects of various factors affecting oxygen delivery/consumption balance on tissue oxygen extraction and to determine specific NIRS-derived FTOE patterns to aid in incorporating this noninvasive monitoring strategy.

STATEMENT OF FINANCIAL SUPPORT

This study was supported in part by an unrestricted equipment use grant from Somanetics (now Medtronic, Boulder, CO).

DISCLOSURE

All authors have no financial relationships or conflicts of interest to disclose related to this study.

ACKNOWLEDGMENTS

We would like to thank the families of our patients for allowing us to perform NIRS monitoring on their VLBW neonates. We also want to acknowledge the cooperation of our nursing and medical staff during the conduct of this work.

REFERENCES

1. Patel RM, Knezevic A, Shenvi N, et al. Association of red blood cell transfusion, anemia, and necrotizing enterocolitis in very low-birth-weight infants. *JAMA* 2016;315:889-97
2. Klein HG, Flegel WA, Natanson C. Red Blood cell transfusion: precision vs imprecision medicine. *JAMA* 2015;314:1557-8
3. Kirpalani H, Whyte RK, Andersen C, et al. The premature infants in need of transfusion (pint) study: a randomized, controlled trial of a restrictive (low) versus liberal (high) transfusion threshold for extremely low birth weight infants. *J Pediatr* 2006;149:301-7
4. Whyte R, Kirpalani H. Low versus high haemoglobin concentration threshold for blood transfusion for preventing morbidity and mortality in very low birth weight infants. *Cochrane Database Syst Rev* 2011;11:CD000512
5. La Gamma EF, Krauss A, Auld PAM. Effects of increased red cell mass on subclinical tissue acidosis in hyaline membrane disease. *Arch Dis Child Fetal Neonatal* Ed 1996;75:F87-93
6. Andersen CC, Hodyl NA, Kirpalani HM, Stark MJ. A theoretical and practical approach to defining "adequate oxygenation" in the preterm newborn. *Pediatrics* 2017;139
7. Sood BG, McLaughlin K, Cortez J. Near-infrared spectroscopy: applications in neonates. *Semin Fetal Neonatal Med* 2015;20:164-72
8. Chakravarti S, Srivastava S, Mitnacht AJC. Near infrared spectroscopy (NIRS) in children. *Semin Cardiothorac Vasc Anesth* 2008;12:70-9

9. Dani C, Pratesi S, Fontanelli G, Barp J, Bertini G. Blood transfusions increase cerebral, splanchnic, and renal oxygenation in anemic preterm infants. *Transfusion* 2010;50:1220-6
10. Mintzer JP, Parvez B, Chelala M, Alpan G, LaGamma EF. Monitoring regional tissue oxygen extraction in neonates <1250 g helps identify transfusion thresholds independent of hematocrit. *J Neonatal Perinatal Med* 2014;7:89-100
11. White L, Said M, Rais-Bahrami K. Monitoring mesenteric tissue oxygenation with near-infrared spectroscopy during packed red blood cell transfusion in preterm infants. *J Neonatal Perinatal Med* 2015;8:157-63
12. Andersen CC, Karayil SM, Hodyl NA, Stark MJ. Early red cell transfusion favourably alters cerebral oxygen extraction in very preterm newborns. *Arch Dis Child Fetal Neonatal Ed* 2015;100:F433-5
13. Seidel D, Blaser A, Gebauer C, Pulzer F, Thome U, Knupfer M. Changes in regional tissue oxygenation saturation and desaturations after red blood cell transfusion in preterm infants. *J Perinatol* 2013;33:282-7
14. van Hoften JCR, Verhagen EA, Keating P, ter Horst HJ, Bos AF. Cerebral tissue oxygen saturation and extraction in preterm infants before and after blood transfusion. *Arch Dis Child Fetal Neonatal Ed* 2010;95:F352-F8
15. Wardle SP, Garr R, Yoxall CW, Weindling AM. A pilot randomised controlled trial of peripheral fractional oxygen extraction to guide blood transfusions in preterm infants. *Arch Dis Child Fetal Neonatal Ed* 2002;86:F22-7

16. Mintzer JP, Parvez B, Chelala M, Alpan G, LaGamma EF. Quiescent variability of cerebral, renal, and splanchnic regional tissue oxygenation in very low birth weight neonates. *J Neonatal Perinatal Med* 2014;7:199-206
17. Mintzer JP, Parvez B, Alpan G, LaGamma EF. Effects of sodium bicarbonate correction of metabolic acidosis on regional tissue oxygenation in very low birth weight neonates. *J Perinatol* 2015;35:601-6
18. Mintzer JP, Parvez B, La Gamma EF. Umbilical arterial blood sampling alters cerebral tissue oxygenation in very low birth weight neonates. *J Pediatr* 2015;167:1013-7
19. Hyttel-Sorensen S, Pellicer A, Alderliesten T, et al. Cerebral near infrared spectroscopy oximetry in extremely preterm infants: phase II randomised clinical trial. *BMJ* 2015;350:g7635
20. Somanetics. Troy, MI; cited 2011 November [about 1 screen]. Available from: <http://www.somanetics.com>
21. McNeill S, Gatenby JC, McElroy S, Engelhardt B. Normal cerebral, renal and abdominal regional oxygen saturations using near-infrared spectroscopy in preterm infants. *J Perinatol* 2010;31:51-7
22. Bland JM, Altman DG. Calculating correlation coefficients with repeated observations: Part 1--Correlation within subjects. *BMJ* 1995;310:446
23. Alkalay AL, Galvis S, Ferry DA, Simmons CF, Krueger RC, Jr. Hemodynamic changes in anemic premature infants: are we allowing the hematocrits to fall too low? *Pediatrics* 2003;112:838-45

24. Bailey S, Hendricks-Muñoz K, Wells J, Mally P. Packed red blood cell transfusion increases regional cerebral and splanchnic tissue oxygen saturation in anemic symptomatic preterm infants. *Am J Perinatol* 2010;27:445-53
25. Sandal G, Oguz SS, Erdevi O, Akar M, Uras N, Dilmen U. Assessment of red blood cell transfusion and transfusion duration on cerebral and mesenteric oxygenation using near-infrared spectroscopy in preterm infants with symptomatic anemia. *Transfusion* 2014;54:1100-5
26. Bailey SM, Hendricks-Munoz KD, Mally PV. Variability in splanchnic tissue oxygenation during preterm red blood cell transfusion given for symptomatic anaemia may reveal a potential mechanism of transfusion-related acute gut injury. *Blood Transfus* 2015;13:429-34
27. Giesinger RE, McNamara PJ. Hemodynamic instability in the critically ill neonate: An approach to cardiovascular support based on disease pathophysiology. *Semin Perinatol* 2016;40:174-88
28. Chock VY, Rose LA, Mante JV, Punj R. Near-infrared spectroscopy for detection of a significant patent ductus arteriosus. *Pediatr Res* 2016;80:675-80
29. Stark MJ, Hodyl NA, Belegar VK, Andersen CC. Intrauterine inflammation, cerebral oxygen consumption and susceptibility to early brain injury in very preterm newborns. *Arch Dis Child Fetal Neonatal Ed* 2016;101:F137-42
30. Akotia DH, Durham JT, Arnell KM, Petruzzelli DL, Katheria AC. Relationship between near-infrared spectroscopy and transabdominal ultrasonography: noninvasive monitoring of intestinal function in neonates. *Med Sci Monit* 2016;22:61-8

FIGURE CAPTIONS

Figure 1: Cerebral and flank fractional tissue oxygen extraction inversely correlated with hematocrit level; an effect not observed for the splanchnic site. Fractional tissue oxygen extraction was smallest and least variable at the cerebral site, intermediate in flank, and highest and most variable at the splanchnic monitoring site.

Figure 2: Corresponding hematocrit values for individual hematocrit quartiles.

** $p < 0.001$ for ANOVA comparisons of all hematocrit quartiles and post-hoc comparisons between Q1/Q2/Q3 and Q4 hematocrit quartiles.

Figure 3: Median and interquartile ranges for cerebral, flank, and splanchnic regional oxygen saturations corresponding to individual hematocrit quartiles. ** $p < 0.001$ for cerebral rSO₂ ANOVA comparison of all hematocrit quartiles and post-hoc comparisons between Q1/Q2/Q3 and Q4 hematocrit quartiles. ++ $p < 0.001$ for flank rSO₂ ANOVA comparison of all hematocrit quartiles and post-hoc comparisons between Q1/Q2 and Q4 hematocrit quartiles. No statistically significant interquartile differences were noted for splanchnic rSO₂.

Figure 4: Median and interquartile ranges for cerebral, flank, and splanchnic fractional tissue oxygen extraction corresponding to individual hematocrit quartiles. ** $p < 0.05$ for cerebral FTOE ANOVA comparison of all hematocrit quartiles and post-hoc comparisons between Q1 ($p = 0.004$), Q2 ($p = 0.01$), and Q3 ($p = 0.03$) and Q4 hematocrit quartiles. ++ $p < 0.001$ for flank FTOE ANOVA comparison of all hematocrit quartiles and post-hoc comparisons between Q1/Q2 and Q4 hematocrit quartiles. No statistically significant interquartile differences were noted for splanchnic FTOE.

TABLES

Table 1: Hematocrits and corresponding vital signs based on hematocrit quartile.

		P-value
Hematocrit (%)		<0.001 (ANOVA)
1 st Quartile	33.3 ± 2.1 (33.8; 27.3-35.8)	<0.001
2 nd Quartile	37.5 ± 1.0 (37.4; 35.9-39.0)	<0.001
3 rd Quartile	40.7 ± 1.0 (40.7; 39.0-42.3)	<0.001
4 th Quartile	46.8 ± 3.2 (46.5; 42.3-52.9)	--
ALL	39.7 ± 5.4 (39.0; 27.3-52.9)	
Heart Rate (beats/min)		0.30 (ANOVA)
1 st Quartile	158 ± 11 (159; 137-182)	NS
2 nd Quartile	158 ± 13 (157; 131-181)	NS
3 rd Quartile	162 ± 10 (161; 136-181)	NS
4 th Quartile	157 ± 10 (157; 138-175)	--
ALL	159 ± 11 (159; 131-182)	
Systolic Blood Pressure (mmHg)		0.005 (ANOVA)
1 st Quartile	50 ± 8 (48; 38-78)	0.003
2 nd Quartile	51 ± 9 (39; 37-74)	0.03
3 rd Quartile	51 ± 10 (48; 37-76)	NS
4 th Quartile	56 ± 8 (58; 37-71)	--
ALL	52 ± 9 (50; 37-78)	
Diastolic Blood Pressure		<0.001 (ANOVA)

(mmHg)		
1 st Quartile	25 ± 7 (24; 17-42)	<0.001
2 nd Quartile	27 ± 7 (27; 17-47)	<0.001
3 rd Quartile	27 ± 7 (27; 13-40)	<0.001
4 th Quartile	34 ± 6 (35; 19-49)	--
ALL	28 ± 7 (28; 13-49)	
Pulse Oximetry (%)		0.013 (ANOVA)
1 st Quartile	93 ± 4 (93; 84-100)	0.006
2 nd Quartile	92 ± 5 (93; 79-100)	0.008
3 rd Quartile	93 ± 5 (94; 79-100)	NS
4 th Quartile	96 ± 4 (97; 85-100)	--
ALL	93 ± 5 (95; 79-100)	

Data expressed as mean ± SD (median; range)

Comparisons via ANOVA (or ANOVA on ranks) with Dunnett (or Dunn's, if ranks used) post-hoc test comparing to 4th hematocrit quartile.

Table 2: Near-infrared spectroscopy data based on hematocrit quartile.

		P-value
Cerebral rSO₂ (%)		<0.001 (ANOVA)
1 st Quartile	69 ± 7 (70; 53-81)	<0.001
2 nd Quartile	71 ± 10 (70; 48-95)	<0.001
3 rd Quartile	70 ± 10 (71; 46-85)	<0.001
4 th Quartile	79 ± 11 (79; 46-93)	--
ALL	72 ± 10 (72; 46-95)	
Flank rSO₂ (%)		<0.001 (ANOVA)
1 st Quartile	59 ± 12 (59; 41-83)	<0.001
2 nd Quartile	61 ± 12 (62; 38-79)	<0.001
3 rd Quartile	64 ± 18 (68; 27-90)	NS
4 th Quartile	75 ± 13 (78; 38-95)	--
ALL	65 ± 15 (66; 27-95)	
Splanchnic rSO₂ (%)		0.10 (ANOVA)
1 st Quartile	36 ± 18 (35; 15-87)	NS
2 nd Quartile	39 ± 18 (36; 15-70)	NS
3 rd Quartile	48 ± 21 (44; 18-91)	NS
4 th Quartile	38 ± 19 (35; 16-77)	--
ALL	40 ± 19 (38; 15-91)	
Cerebral FTOE		0.004 (ANOVA)
1 st Quartile	0.26 ± 0.08 (0.25; 0.10-0.41)	0.004
2 nd Quartile	0.24 ± 0.09 (0.25; 0.04-0.45)	0.01
3 rd Quartile	0.25 ± 0.09 (0.23; 0.12-0.42)	0.03

4 th Quartile	0.18 ± 0.10 (0.18; 0.04-0.46)	--
ALL	0.23 ± 0.10 (0.22; 0.04-0.46)	
Flank FTOE		<0.001 (ANOVA)
1 st Quartile	0.36 ± 0.12 (0.38; 0.14-0.56)	<0.001
2 nd Quartile	0.35 ± 0.11 (0.35; 0.20-0.56)	<0.001
3 rd Quartile	0.32 ± 0.17 (0.29; 0.09-0.68)	NS
4 th Quartile	0.22 ± 0.13 (0.19; 0.03-0.59)	--
ALL	0.31 ± 0.14 (0.30; 0.03-0.68)	
Splanchnic FTOE		0.18 (ANOVA)
1 st Quartile	0.61 ± 0.20 (0.64; 0.06-0.84)	NS
2 nd Quartile	0.58 ± 0.21 (0.60; 0.16-0.85)	NS
3 rd Quartile	0.50 ± 0.21 (0.52; 0.04-0.80)	NS
4 th Quartile	0.60 ± 0.20 (0.64; 0.20-0.84)	--
ALL	0.57 ± 0.21 (0.59; 0.04-0.85)	

Data expressed as mean ± SD (median; range).

Comparisons via ANOVA (or ANOVA on ranks) with Dunnett (or Dunn's, if ranks used) post-hoc test comparing to 4th hematocrit quartile.

Table 3: Multivariate logistic regression including hypotension requiring inotropic support; hemodynamically-significant patent ductus arteriosus; and intraventricular

hemorrhage. Hct = hematocrit; Dopa = dopamine administration; PDA =

hemodynamically-significant patent ductus arteriosus; IVH = intraventricular

hemorrhage.

Site	Factor	R-Square Change	P-value
Cerebral FTOE	Hct	0.136	<0.0001
	Hct	0.136	<0.0001
	Dopa	0.115	<0.0001
	Hct	0.136	<0.0001
	PDA	0.006	0.39
	Hct	0.136	<0.0001
	IVH	0.013	0.21
	Hct	0.136	0.001
	Dopa	0.115	<0.0001
	PDA	0.002	0.61
	IVH	0.035	0.02
Flank FTOE	Hct	0.176	<0.0001
	Hct	0.176	<0.0001
	Dopa	0.158	<0.0001
	Hct	0.176	<0.0001
	PDA	0.122	<0.0001
	Hct	0.176	<0.0001
	IVH	0.002	0.62
	Hct	0.176	0.001
	Dopa	0.158	<0.0001
	PDA	0.045	0.007
	IVH	0.004	0.43
Splanchnic FTOE	Hct	0.011	0.28
	Hct	0.011	0.19
	Dopa	0.021	0.13
	Hct	0.011	0.02
	PDA	0.148	<0.0001

	Hct	0.011	0.48
	IVH	0.038	0.04
	Hct	0.011	0.04
	Dopa	0.021	0.84
	PDA	0.127	<0.0001
	IVH	0.023	0.09